

Question ID 92921647

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: 92921647

For the function f , the graph of $y = f(x)$ in the xy -plane has a slope of 3 and passes through the point $(0, -8)$. Which equation defines f ?

- A. $f(x) = 3x$
- B. $f(x) = 3x - 8$
- C. $f(x) = 3x + 5$
- D. $f(x) = 3x + 11$

ID: 92921647 Answer

Correct Answer: B

Rationale

Choice B is correct. An equation defining a linear function can be written in the form $f(x) = mx + b$, where m and b are constants, m is the slope of the graph of $y = f(x)$ in the xy -plane, and $(0, b)$ is the y -intercept of the graph. It's given that for the function f , the graph of $y = f(x)$ in the xy -plane has a slope of 3 . Therefore, $m = 3$. It's also given that this graph passes through the point $(0, -8)$. Therefore, the y -intercept of the graph is $(0, -8)$, and it follows that $b = -8$. Substituting 3 for m and -8 for b in the equation $f(x) = mx + b$ yields $f(x) = 3x - 8$. Thus, the equation that defines f is $f(x) = 3x - 8$.

Choice A is incorrect. For this function, the graph of $y = f(x)$ in the xy -plane passes through the point $(0, 0)$, not $(0, -8)$.

Choice C is incorrect. For this function, the graph of $y = f(x)$ in the xy -plane passes through the point $(0, 5)$, not $(0, -8)$.

Choice D is incorrect. For this function, the graph of $y = f(x)$ in the xy -plane passes through the point $(0, 11)$, not $(0, -8)$.

Question Difficulty: Easy

Question ID e1f981b8

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: e1f981b8

The function g is defined by $g(x) = 6x$. For what value of x is $g(x) = 54$?

ID: e1f981b8 Answer

Correct Answer: 9

Rationale

The correct answer is **9**. It's given that $g(x) = 6x$. Substituting **54** for $g(x)$ in the given function yields $54 = 6x$. Dividing both sides of this equation by **6** yields $x = 9$. Therefore, the value of x when $g(x) = 54$ is **9**.


Question Difficulty: Easy

Question ID 3eef8a08

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: 3eef8a08

Hydrogen is placed inside a container and kept at a constant pressure. The graph shows the estimated volume y , in liters, of the hydrogen when its temperature is x kelvins.



What is the estimated volume, in liters, of the hydrogen when its temperature is **500** kelvins?

- A. 0
- B. $\frac{7}{500}$
- C. 7
- D. $\frac{500}{7}$

ID: 3eef8a08 Answer

Correct Answer: C

Rationale

Choice C is correct. For the graph shown, the x -axis represents temperature, in kelvins, and the y -axis represents volume, in liters. Therefore, the estimated volume, in liters, of the hydrogen when its temperature is **500** kelvins is represented by the y -coordinate of the point on the graph that has an x -coordinate of **500**. The point on the graph with an x -coordinate of **500** has a y -coordinate of **7**. Therefore, the estimated volume, in liters, of the hydrogen when its temperature is **500** kelvins is **7**.

Choice A is incorrect and may result from conceptual errors.

Choice B is incorrect and may result from conceptual errors.

Choice D is incorrect and may result from conceptual errors.

Question Difficulty: Easy

Question ID 26e3a069

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: 26e3a069

The function g is defined by $g(x) = 10x + 8$. What is the value of $g(x)$ when $x = 8$?

- A. 0
- B. 8
- C. 10
- D. 88

ID: 26e3a069 Answer

Correct Answer: D

Rationale

Choice D is correct. The value of $g(x)$ when $x = 8$ can be found by substituting 8 for x in the given equation $g(x) = 10x + 8$. This yields $g(8) = 10(8) + 8$, or $g(8) = 88$. Therefore, when $x = 8$, the value of $g(x)$ is 88.

Choice A is incorrect. This is the value of x when $g(x) = 8$, rather than the value of $g(x)$ when $x = 8$.

Choice B is incorrect and may result from conceptual or calculation errors.

Choice C is incorrect and may result from conceptual or calculation errors.

Question Difficulty: Easy

Question ID bc1e9b9b

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: bc1e9b9b

x	$f(x)$
0	29
1	32
2	35

For the linear function f , the table shows three values of x and their corresponding values of $f(x)$. Which equation defines $f(x)$?

- A. $f(x) = 3x + 29$
- B. $f(x) = 29x + 32$
- C. $f(x) = 35x + 29$
- D. $f(x) = 32x + 35$

ID: bc1e9b9b Answer

Correct Answer: A

Rationale

Choice A is correct. An equation that defines a linear function f can be written in the form $f(x) = mx + b$, where m and b are constants. It's given in the table that when $x = 0$, $f(x) = 29$. Substituting 0 for x and 29 for $f(x)$ in the equation $f(x) = mx + b$ yields $29 = m(0) + b$, or $29 = b$. Substituting 29 for b in the equation $f(x) = mx + b$ yields $f(x) = mx + 29$. It's also given in the table that when $x = 1$, $f(x) = 32$. Substituting 1 for x and 32 for $f(x)$ in the equation $f(x) = mx + 29$ yields $32 = m(1) + 29$, or $32 = m + 29$. Subtracting 29 from both sides of this equation yields $3 = m$. Substituting 3 for m in the equation $f(x) = mx + 29$ yields $f(x) = 3x + 29$.

Choice B is incorrect and may result from conceptual or calculation errors.

Choice C is incorrect and may result from conceptual or calculation errors.

Choice D is incorrect and may result from conceptual or calculation errors.

Question Difficulty: Easy

Question ID 389861fc

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: 389861fc

The function h is defined by $h(x) = x + 200$. What is the value of $h(50)$?

- A. 200
- B. 250
- C. 10,000
- D. 50,200

ID: 389861fc Answer

Correct Answer: B

Rationale

Choice B is correct. Substituting 50 for x in the given function yields $h(50) = 50 + 200$, or $h(50) = 250$. Therefore, the value of $h(50)$ is 250 .

Choice A is incorrect. This is the value of $h(0)$.

Choice C is incorrect. This is the value of $h(9,800)$.

Choice D is incorrect. This is the value of $h(50,000)$.

Question Difficulty: Easy

Question ID 76f44a77

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: 76f44a77

The function f is defined by $f(x) = 8x$. For what value of x does $f(x) = 72$?

- A. 8
- B. 9
- C. 64
- D. 80

ID: 76f44a77 Answer

Correct Answer: B

Rationale

Choice B is correct. Substituting 72 for $f(x)$ in the given function yields $72 = 8x$. Dividing each side of this equation by 8 yields $9 = x$. Therefore, $f(x) = 72$ when the value of x is 9 .

Choice A is incorrect. This is the value of x for which $f(x) = 64$, not $f(x) = 72$.

Choice C is incorrect. This is the value of x for which $f(x) = 512$, not $f(x) = 72$.

Choice D is incorrect. This is the value of x for which $f(x) = 640$, not $f(x) = 72$.

Question Difficulty: Easy

Question ID 4c580285

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: 4c580285

The function f is defined by the equation $f(x) = 7x + 2$. What is the value of $f(x)$ when $x = 4$?

ID: 4c580285 Answer

Correct Answer: 30

Rationale

The correct answer is **30**. The value of $f(x)$ when $x = 4$ can be found by substituting **4** for x in the given equation $f(x) = 7x + 2$. This yields $f(4) = 7(4) + 2$, or $f(4) = 30$. Therefore, when $x = 4$, the value of $f(x)$ is **30**.

Question Difficulty: Easy

Question ID dc85c551

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: dc85c551

$$d = 16t$$

The given equation represents the distance d , in inches, where t represents the number of seconds since an object started moving. Which of the following is the best interpretation of 16 in this context?

- A. The object moved a total of 16 inches.
- B. The object moved a total of $16t$ inches.
- C. The object is moving at a rate of 16 inches per second.
- D. The object is moving at a rate of $\frac{1}{16}$ inches per second.

ID: dc85c551 Answer

Correct Answer: C

Rationale

Choice C is correct. It's given that in the equation $d = 16t$, d represents the distance, in inches, and t represents the number of seconds since an object started moving. In this equation, t is being multiplied by 16 . This means that the object's distance increases by 16 inches each second. Therefore, the best interpretation of 16 in this context is that the object is moving at a rate of 16 inches per second.

Choice A is incorrect and may result from conceptual errors.

Choice B is incorrect. This is the best interpretation of $16t$, rather than 16 , in this context.

Choice D is incorrect and may result from conceptual errors.

Question Difficulty: Easy

Question ID 15bcfa0b

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: 15bcfa0b

The function f is defined by $f(x) = 4x$. For what value of x does $f(x) = 8$?

ID: 15bcfa0b Answer

Correct Answer: 2

Rationale

The correct answer is **2**. Substituting **8** for $f(x)$ in the given equation yields $8 = 4x$. Dividing the left- and right-hand sides of this equation by **4** yields $x = 2$. Therefore, the value of x is **2** when $f(x) = 8$.

Question Difficulty: Easy

Question ID 5eeef20e

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: 5eeef20e

A veterinarian recommends that each day a certain rabbit should eat **25** calories per pound of the rabbit’s weight, plus an additional **11** calories. Which equation represents this situation, where *c* is the total number of calories the veterinarian recommends the rabbit should eat each day if the rabbit’s weight is *x* pounds?

- A. $c = 25x$
- B. $c = 36x$
- C. $c = 11x + 25$
- D. $c = 25x + 11$

ID: 5eeef20e Answer

Correct Answer: D

Rationale

Choice D is correct. It’s given that a veterinarian recommends that each day the rabbit should eat **25** calories per pound of the rabbit’s weight, plus an additional **11** calories. If the rabbit’s weight is *x* pounds, then multiplying **25** calories per pound by the rabbit’s weight, *x* pounds, yields **25*x*** calories. Adding the additional **11** calories that the rabbit should eat each day yields **25*x* + 11** calories. It’s given that *c* is the total number of calories the veterinarian recommends the rabbit should eat each day if the rabbit’s weight is *x* pounds. Therefore, this situation can be represented by the equation $c = 25x + 11$.

Choice A is incorrect. This equation represents a situation where a veterinarian recommends that each day the rabbit should eat **25** calories per pound of the rabbit’s weight.

Choice B is incorrect. This equation represents a situation where a veterinarian recommends that each day the rabbit should eat **25 + 11**, or **36**, calories per pound of the rabbit’s weight.

Choice C is incorrect. This equation represents a situation where a veterinarian recommends that each day the rabbit should eat **11** calories per pound of the rabbit’s weight, plus an additional **25** calories.


Question Difficulty: Easy

Question ID b15c634e

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: b15c634e

The graph shown models the number of candy bars a certain machine wraps with a label in x seconds.



According to the graph, what is the estimated number of candy bars the machine wraps with a label per second?

- A. 2
- B. 40
- C. 78
- D. 80

ID: b15c634e Answer

Correct Answer: B

Rationale

Choice B is correct. For the graph shown, the x -axis represents time, in seconds, and the y -axis represents the number of candy bars wrapped. The slope of a line in the xy -plane is the change in y for each 1-unit increase in x . It follows that the slope of the graph shown represents the estimated number of candy bars the machine wraps with a label per second. The slope, m , of a line in the xy -plane can be found using any two points, (x_1, y_1) and (x_2, y_2) , on the line and the slope formula $m = \frac{y_2 - y_1}{x_2 - x_1}$. The graph shown passes through the points $(0, 0)$ and $(2, 80)$. Substituting these points for (x_1, y_1) and (x_2, y_2) , respectively, in the slope formula yields $m = \frac{80 - 0}{2 - 0}$, which is equivalent to $m = \frac{80}{2}$, or $m = 40$. Therefore, the estimated number of candy bars the machine wraps with a label per second is 40.

Choice A is incorrect and may result from conceptual or calculation errors.

Choice C is incorrect and may result from conceptual or calculation errors.

Choice D is incorrect and may result from conceptual or calculation errors.


Question Difficulty: Easy

Question ID a4e1df2e

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: a4e1df2e

The cost y , in dollars, for a manufacturer to make x rings is represented by the line shown.



What is the cost, in dollars, for the manufacturer to make **60** rings?

- A. 100
- B. 125
- C. 175
- D. 225

ID: a4e1df2e Answer

Correct Answer: C

Rationale

Choice C is correct. The line shown represents the cost y , in dollars, for a manufacturer to make x rings. For the line shown, the x -axis represents the number of rings made by the manufacturer and the y -axis represents the cost, in dollars. Therefore, the cost, in dollars, for the manufacturer to make **60** rings is represented by the y -coordinate of the point on the line that has an x -coordinate of **60**. The point on the line with an x -coordinate of **60** has a y -coordinate of **175**. Therefore, the cost, in dollars, for the manufacturer to make **60** rings is **175**.

Choice A is incorrect and may result from conceptual or calculation errors.

Choice B is incorrect. This is the cost, in dollars, for the manufacturer to make **20** rings.


Choice D is incorrect. This is the cost, in dollars, for the manufacturer to make **100** rings.

Question Difficulty: Easy

Question ID 7d9d3fb4

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: 7d9d3fb4



The graph of the linear function f is shown, where $y = f(x)$. What is the x-intercept of the graph of f ?

- A. $(-12, 0)$
- B. $(0, 0)$
- C. $(\frac{1}{4}, 0)$
- D. $(12, 0)$

ID: 7d9d3fb4 Answer

Correct Answer: A

Rationale

Choice A is correct. The x-intercept of a graph is the point where the graph intersects the x-axis. The graph of function f , where $y = f(x)$, intersects the x-axis at $(-12, 0)$. Therefore, the x-intercept of the graph of f is $(-12, 0)$.

Choice B is incorrect and may result from conceptual or calculation errors.

Choice C is incorrect and may result from conceptual or calculation errors.

Choice D is incorrect and may result from conceptual or calculation errors.

Question Difficulty: Easy

Question ID 0e2e504f

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: 0e2e504f

The function f is defined by $f(x) = \frac{1}{10}x - 2$. What is the y -intercept of the graph of $y = f(x)$ in the xy -plane?

- A. $(-2, 0)$
- B. $(0, -2)$
- C. $(0, \frac{1}{10})$
- D. $(\frac{1}{10}, 0)$

ID: 0e2e504f Answer

Correct Answer: B

Rationale

Choice B is correct. The y -intercept of the graph of a function in the xy -plane is the point on the graph where $x = 0$. It's given that $f(x) = \frac{1}{10}x - 2$. Substituting 0 for x in this equation yields $f(0) = \frac{1}{10}(0) - 2$, or $f(0) = -2$. Since it's given that $y = f(x)$, it follows that $y = -2$ when $x = 0$. Therefore, the y -intercept of the graph of $y = f(x)$ in the xy -plane is $(0, -2)$.

Choice A is incorrect and may result from conceptual or calculation errors.

Choice C is incorrect and may result from conceptual or calculation errors.

Choice D is incorrect and may result from conceptual or calculation errors.

Question Difficulty: Easy

Question ID 3c80bb57

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: 3c80bb57

The total cost $f(x)$, in dollars, to lease a car for **36** months from a particular car dealership is given by $f(x) = 36x + 1,000$, where x is the monthly payment, in dollars. What is the total cost to lease a car when the monthly payment is **\$400**?

- A. **\$13,400**
- B. **\$13,000**
- C. **\$15,400**
- D. **\$37,400**

ID: 3c80bb57 Answer

Correct Answer: C

Rationale

Choice C is correct. It's given that $f(x)$ is the total cost, in dollars, to lease a car from this dealership with a monthly payment of x dollars. Therefore, the total cost, in dollars, to lease the car when the monthly payment is **\$400** is represented by the value of $f(x)$ when $x = 400$. Substituting **400** for x in the equation $f(x) = 36x + 1,000$ yields $f(400) = 36(400) + 1,000$, or $f(400) = 15,400$. Thus, when the monthly payment is **\$400**, the total cost to lease a car is **\$15,400**.

Choice A is incorrect and may result from conceptual or calculation errors.

Choice B is incorrect and may result from conceptual or calculation errors.

Choice D is incorrect and may result from conceptual or calculation errors.

Question Difficulty: Easy

Question ID 61ab5e49

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: 61ab5e49

The function f is defined by $f(x) = 80 - 6x$. What is the value of $f(7)$?

- A. 13
- B. 38
- C. 74
- D. 81

ID: 61ab5e49 Answer

Correct Answer: B

Rationale

Choice B is correct. It's given that function f is defined by $f(x) = 80 - 6x$. The value of $f(7)$ can be found by substituting 7 for x in the given function, which yields $f(7) = 80 - 6(7)$, or $f(7) = 80 - 42$, which is equivalent to $f(7) = 38$. Therefore, the value of $f(7)$ is 38.

Choice A is incorrect. This is the value of $80 - 67$, not $80 - 6(7)$.

Choice C is incorrect. This is the value of $80 - 6(1)$, not $80 - 6(7)$.

Choice D is incorrect. This is the value of $80 - 6 + 7$, not $80 - 6(7)$.

Question Difficulty: Easy

Question ID 4453d265

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: 4453d265

The number y is ~~84~~ less than the number x . Which equation represents the relationship between x and y ?

- A. $y = x + 84$
- B. $y = \frac{1}{84}x$
- C. $y = 84x$
- D. $y = x - 84$

ID: 4453d265 Answer

Correct Answer: D

Rationale

Choice D is correct. It’s given that the number y is ~~84~~ less than the number x . A number that's ~~84~~ less than the number x is equivalent to ~~84~~ subtracted from the number x , or $x - \del{84}$. Therefore, the equation $y = x - 84$ represents the relationship between x and y .

Choice A is incorrect and may result from conceptual errors.

Choice B is incorrect and may result from conceptual errors.


Choice C is incorrect and may result from conceptual errors.

Question Difficulty: Easy

Question ID 99235d83

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: 99235d83



The graph of the linear function f is shown, where $y = f(x)$. What is the y -intercept of the graph of f ?

- A. $(0, 0)$
- B. $(0, -\frac{16}{11})$
- C. $(0, -8)$
- D. $(0, 8)$

ID: 99235d83 Answer

Correct Answer: D

Rationale

Choice D is correct. The y -intercept of a graph is the point where the graph intersects the y -axis. The graph of function f shown intersects the y -axis at the point $(0, 8)$. Therefore, the y -intercept of the graph of f is $(0, 8)$.

Choice A is incorrect. This is the point where the x -axis, not the graph of f , intersects the y -axis.

Choice B is incorrect and may result from conceptual or calculation errors.

Choice C is incorrect and may result from conceptual or calculation errors.

Question Difficulty: Easy

Question ID 87bd4e09

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: 87bd4e09

The function f is defined by the equation $f(x) = 100x + 2$. What is the value of $f(x)$ when $x = 9$?

- A. 111
- B. 118
- C. 900
- D. 902

ID: 87bd4e09 Answer

Correct Answer: D

Rationale

Choice D is correct. Substituting 9 for x in the given equation yields $f(9) = 100(9) + 2$, or $f(9) = 902$. Therefore, the value of $f(x)$ when $x = 9$ is 902.

Choice A is incorrect. This is the value of $f(x)$ when $x = 1.09$.

Choice B is incorrect. This is the value of $f(x)$ when $x = 1.16$.

Choice C is incorrect. This is the value of $f(x)$ when $x = 8.98$.

Question Difficulty: Easy

Question ID 6bd32629

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: 6bd32629

$$f(x) = 7x + 1$$

The function gives the total number of people on a company retreat with x managers. What is the total number of people on a company retreat with 7 managers?

ID: 6bd32629 Answer

Correct Answer: 50

Rationale

The correct answer is **50**. It's given that the function f gives the total number of people on a company retreat with x managers. It's also given that 7 managers are on the company retreat. Substituting 7 for x in the given function yields $f(7) = 7(7) + 1$, or $f(7) = 50$. Therefore, there are a total of **50** people on a company retreat with 7 managers.

Question Difficulty: Easy

Question ID 6a8610a5

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: 6a8610a5

If $y = 5x + 10$, what is the value of y when $x = 8$?

ID: 6a8610a5 Answer

Correct Answer: 50

Rationale

The correct answer is **50**. Substituting **8** for x in the given equation yields $y = 5(8) + 10$, or $y = 50$. Therefore, the value of y is **50** when $x = 8$.


Question Difficulty: Easy

Question ID 8b896724

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: 8b896724

Oxygen gas is placed inside a tank with a constant volume. The graph shows the estimated temperature y , in kelvins, of the oxygen gas when its pressure is x atmospheres.



What is the estimated temperature, in kelvins, of the oxygen gas when its pressure is **6** atmospheres?

- A. **6**
- B. **60**
- C. **700**
- D. **760**

ID: 8b896724 Answer

Correct Answer: C

Rationale

Choice C is correct. For the graph shown, the x -axis represents pressure, in atmospheres, and the y -axis represents temperature, in kelvins. Therefore, the estimated temperature, in kelvins, of the oxygen gas when its pressure is **6** atmospheres is represented by the y -coordinate of the point on the graph that has an x -coordinate of **6**. The point on the graph with an x -coordinate of **6** has a y -coordinate of approximately **700**. Therefore, the estimated temperature, in kelvins, of the oxygen gas when its pressure is **6** atmospheres is **700**.

Choice A is incorrect. This is the pressure, in atmospheres, not the estimated temperature, in kelvins, of the oxygen gas when its pressure is **6** atmospheres.

Choice B is incorrect and may result from conceptual or calculation errors.

Choice D is incorrect and may result from conceptual or calculation errors.

Question Difficulty: Easy

Question ID 273c9f49

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: 273c9f49

The function h is defined by $h(x) = 3x - 7$. What is the value of $h(-2)$?

- A. -13
- B. -10
- C. 10
- D. 13

ID: 273c9f49 Answer

Correct Answer: A

Rationale

Choice A is correct. The value of $h(-2)$ can be found by substituting -2 for x in the equation defining h . Substituting -2 for x in $h(x) = 3x - 7$ yields $h(-2) = 3(-2) - 7$, or $h(-2) = -13$. Therefore, the value of $h(-2)$ is -13 .

Choice B is incorrect. This is the value of $h(-1)$, not $h(-2)$.

Choice C is incorrect and may result from conceptual or calculation errors.

Choice D is incorrect and may result from conceptual or calculation errors.

Question Difficulty: Easy

Question ID 675325b9

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: 675325b9

Hana deposited a fixed amount into her bank account each month. The function $f(t) = 100 + 25t$ gives the amount, in dollars, in Hana's bank account after t monthly deposits. What is the best interpretation of **25** in this context?

- A. With each monthly deposit, the amount in Hana's bank account increased by **\$25**.
- B. Before Hana made any monthly deposits, the amount in her bank account was **\$25**.
- C. After **1** monthly deposit, the amount in Hana's bank account was **\$25**.
- D. Hana made a total of **25** monthly deposits.

ID: 675325b9 Answer

Correct Answer: A

Rationale

Choice A is correct. It's given that t represents the number of monthly deposits. In the given function $f(t) = 100 + 25t$, the coefficient of t is **25**. This means that for every increase in the value of t by **1**, the value of $f(t)$ increases by **25**. It follows that with each monthly deposit, the amount in Hana's bank account increased by **\$25**.

Choice B is incorrect. Before Hana made any monthly deposits, the amount in her bank account was **\$100**.

Choice C is incorrect. After **1** monthly deposit, the amount in Hana's bank account was **\$125**.


Choice D is incorrect and may result from conceptual errors.

Question Difficulty: Easy

Question ID f717089b

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: f717089b



The graph of the function f is shown, where $y = f(x)$. What is the y -intercept of the graph?

- A. $(0, -1)$
- B. $(0, -4)$
- C. $(0, 1)$
- D. $(0, 4)$

ID: f717089b Answer

Correct Answer: B

Rationale

Choice B is correct. The y -intercept of a graph is the point where the graph intersects the y -axis. The graph of function f shown intersects the y -axis at the point $(0, -4)$. Therefore, the y -intercept of the graph is $(0, -4)$.

Choice A is incorrect and may result from conceptual or calculation errors.

Choice C is incorrect and may result from conceptual or calculation errors.

Choice D is incorrect and may result from conceptual or calculation errors.

Question Difficulty: Easy

Question ID f66cba39

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: f66cba39

Sean rents a tent at a cost of **\$11** per day plus a onetime insurance fee of **\$10**. Which equation represents the total cost c , in dollars, to rent the tent with insurance for d days?

- A. $c = 11(d + 10)$
- B. $c = 10(d + 11)$
- C. $c = 11d + 10$
- D. $c = 10d + 11$

ID: f66cba39 Answer

Correct Answer: C

Rationale

Choice C is correct. It’s given that the cost of renting a tent is **\$11** per day for d days. Multiplying the rental cost by the number of days yields **\$11 d** , which represents the cost of renting the tent for d days before the insurance is added. Adding the onetime insurance fee of **\$10** to the rental cost of **\$11 d** gives the total cost c , in dollars, which can be represented by the equation $c = 11d + 10$.

Choice A is incorrect. This equation represents the total cost to rent the tent if the insurance fee was charged every day.

Choice B is incorrect. This equation represents the total cost to rent the tent if the daily fee was **\$($d + 11$)** for **10** days.


Choice D is incorrect. This equation represents the total cost to rent the tent if the daily fee was **\$10** and the onetime fee was **\$11**.

Question Difficulty: Easy

Question ID 1e0a38e7

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: 1e0a38e7



The graph of the linear function f is shown. What is the y -intercept of the graph of $y = f(x)$?

- A. $(-5, 0)$
- B. $(2, 0)$
- C. $(0, 2)$
- D. $(0, -5)$

ID: 1e0a38e7 Answer

Correct Answer: C

Rationale

Choice C is correct. The y -intercept of a graph is the point where the graph intersects the y -axis. The graph of $y = f(x)$ shown intersects the y -axis at the point $(0, 2)$. Therefore, the y -intercept of the graph of $y = f(x)$ is $(0, 2)$.

Choice A is incorrect. This is the x -intercept, not the y -intercept, of the graph of $y = f(x)$.

Choice B is incorrect and may result from conceptual or calculation errors.

Choice D is incorrect and may result from conceptual or calculation errors.

Question Difficulty: Easy

Question ID afcbd26d

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: afcbd26d

The length, y , of a white whale was **162 centimeters (cm)** when it was born and increased an average of **4.8 cm** per month for the first **12** months after it was born. Which equation best represents this situation, where x is the number of months after the whale was born and y is the length, in **cm**, of the whale?

- A. $y = 162x$
- B. $y = 162x + 162$
- C. $y = 4.8x + 4.8$
- D. $y = 4.8x + 162$

ID: afcbd26d Answer

Correct Answer: D

Rationale

Choice D is correct. It's given that the length of the whale was **162 cm** when it was born and that its length increased an average of **4.8 cm** per month for the first **12** months after it was born. Since x represents the number of months after the whale was born, the total increase in the whale's length, in **cm**, is **4.8** times x , or **4.8x**. The length of the whale y , in **cm**, can be found by adding the whale's length at birth, **162 cm**, to the total increase in length, **4.8x cm**. Therefore, the equation that best represents this situation is $y = 4.8x + 162$.

Choice A is incorrect and may result from conceptual errors.

Choice B is incorrect and may result from conceptual errors.

Choice C is incorrect and may result from conceptual errors.

Question Difficulty: Easy

Question ID 13b10306

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: 13b10306

$$d = 16 - \frac{x}{30}$$

The equation shown gives the estimated amount of diesel d , in gallons, that remains in the gas tank of a truck after being driven x miles, where $0 \leq x \leq 480$. What is the estimated amount of diesel, in gallons, that remains in the gas tank of the truck when $x = 300$?

- A. 0
- B. 6
- C. 14
- D. 16

ID: 13b10306 Answer

Correct Answer: B

Rationale

Choice B is correct. It's given that the equation $d = 16 - \frac{x}{30}$ gives the estimated amount of diesel d , in gallons, that remains in the gas tank of the truck after being driven x miles. Substituting 300 for x in the given equation yields $d = 16 - \frac{300}{30}$, which is equivalent to $d = 16 - 10$, or $d = 6$. Therefore, the estimated amount of diesel that remains in the gas tank of the truck when $x = 300$ is 6 gallons.

Choice A is incorrect. This is the estimated amount of diesel, in gallons, that will remain in the gas tank of the truck when $x = 480$, not when $x = 300$.

Choice C is incorrect. This is the estimated amount of diesel, in gallons, that will remain in the gas tank of the truck when $x = 60$, not when $x = 300$.

Choice D is incorrect. This is the estimated amount of diesel, in gallons, that will remain in the gas tank of the truck when $x = 0$, not when $x = 300$.

Question Difficulty: Easy

Question ID d60b089f

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: d60b089f

The function f is defined by $f(x) = 3x - 8$. What is the value of $f(7)$?

- A. 29
- B. 13
- C. -5
- D. -29

ID: d60b089f Answer

Correct Answer: B

Rationale

Choice B is correct. It's given that the function f is defined by $f(x) = 3x - 8$. The value of $f(7)$ is the value of $f(x)$ when $x = 7$. Substituting 7 for x in the given equation yields $f(7) = 3(7) - 8$, which is equivalent to $f(7) = 21 - 8$, or $f(7) = 13$.

Choice A is incorrect. This is the value of $f(7)$ when $f(x) = 3x + 8$, rather than $f(x) = 3x - 8$.

Choice C is incorrect. This is the value of $f(1)$, rather than $f(7)$.

Choice D is incorrect. This is the value of $f(-7)$, rather than $f(7)$.

Question Difficulty: Easy

Question ID df59edf3

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: df59edf3

A bus is traveling at a constant speed along a straight portion of road. The equation $d = 30t$ gives the distance d , in feet from a road marker, that the bus will be t seconds after passing the marker. How many feet from the marker will the bus be **2** seconds after passing the marker?

- A. **30**
- B. **32**
- C. **60**
- D. **90**

ID: df59edf3 Answer

Correct Answer: C

Rationale

Choice C is correct. It’s given that t represents the number of seconds after the bus passes the marker. Substituting **2** for t in the given equation $d = 30t$ yields $d = 30(2)$, or $d = 60$. Therefore, the bus will be **60** feet from the marker **2** seconds after passing it.

Choice A is incorrect. This is the distance, in feet, the bus will be from the marker **1** second, not **2** seconds, after passing it.

Choice B is incorrect and may result from conceptual or calculation errors.

Choice D is incorrect. This is the distance, in feet, the bus will be from the marker **3** seconds, not **2** seconds, after passing it.

Question Difficulty: Easy

Question ID d76c37db

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: d76c37db

For a training program, Juan rides his bike at an average rate of **5.7** minutes per mile. Which function *m* models the number of minutes it will take Juan to ride *x* miles at this rate?

- A. $m(x) = \frac{x}{5.7}$
- B. $m(x) = x + 5.7$
- C. $m(x) = x - 5.7$
- D. $m(x) = 5.7x$

ID: d76c37db Answer

Correct Answer: D

Rationale

Choice D is correct. It's given that Juan rides his bike at an average rate of **5.7** minutes per mile. The number of minutes it will take Juan to ride *x* miles can be determined by multiplying his average rate by the number of miles, *x*, which yields **5.7x**. Therefore, the function $m(x) = 5.7x$ models the number of minutes it will take Juan to ride *x* miles.

Choice A is incorrect and may result from conceptual errors.

Choice B is incorrect and may result from conceptual errors.

Choice C is incorrect and may result from conceptual errors.

Question Difficulty: Easy

Question ID 5ebcd502

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: 5ebcd502

A contract for a certain service requires a onetime activation cost of **\$35** and a monthly cost of **\$23**. Which equation represents this situation, where c is the total cost, in dollars, of this service contract for t months?

- A. $c = \frac{t}{23} + 35$
- B. $c = \frac{t}{35} + 23$
- C. $c = 23t + 35$
- D. $c = 35t + 23$

ID: 5ebcd502 Answer

Correct Answer: C

Rationale

Choice C is correct. It's given that this service contract requires a monthly cost of **\$23**. A monthly cost of **\$23** for t months results in a cost of **$23t$** . It's also given that this service contract requires a onetime activation cost of **\$35**. Adding the onetime activation cost to the monthly cost of the service contract for t months yields the total cost c , in dollars, of this service contract for t months. Therefore, this situation can be represented by the equation $c = 23t + 35$.

Choice A is incorrect and may result from conceptual or calculation errors.

Choice B is incorrect and may result from conceptual or calculation errors.

Choice D is incorrect and may result from conceptual or calculation errors.


Question Difficulty: Easy

Question ID fa6dad4f

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: fa6dad4f

Argon is placed inside a container with a constant volume. The graph shows the estimated pressure y , in **pounds per square inch (psi)**, of the argon when its temperature is x kelvins.



What is the estimated pressure of the argon, in **psi**, when the temperature is **600** kelvins?

- A. 6
- B. 12
- C. 300
- D. 600

ID: fa6dad4f Answer

Correct Answer: B

Rationale

Choice B is correct. For the graph shown, the x -axis represents temperature, in kelvins, and the y -axis represents the estimated pressure, in **pounds per square inch (psi)**. The estimated pressure of the argon when the temperature is **600** kelvins can be found by locating the point on the graph where the value of x is equal to **600**. The graph passes through the point **(600, 12)**. This means that when the temperature is **600** kelvins, the estimated pressure is **12 psi**.

Choice A is incorrect. This is the estimated pressure, in **psi**, of the argon when the temperature is **300** kelvins, not **600** kelvins.

Choice C is incorrect and may result from conceptual or calculation errors.

Choice D is incorrect. This is the temperature, in kelvins, of the argon.

Question Difficulty: Easy

Question ID 552ce561

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: 552ce561

$$s = 40 + 3t$$

The equation gives the speed s , in miles per hour, of a certain car t seconds after it began to accelerate. What is the speed, in miles per hour, of the car 5 seconds after it began to accelerate?

- A. 40
- B. 43
- C. 45
- D. 55

ID: 552ce561 Answer

Correct Answer: D

Rationale

Choice D is correct. In the given equation, s is the speed, in miles per hour, of a certain car t seconds after it began to accelerate. Therefore, the speed of the car, in miles per hour, 5 seconds after it began to accelerate can be found by substituting 5 for t in the given equation, which yields $s = 40 + 3(5)$, or $s = 55$. Thus, the speed of the car 5 seconds after it began to accelerate is 55 miles per hour.

Choice A is incorrect and may result from conceptual or calculation errors.

Choice B is incorrect and may result from conceptual or calculation errors.

Choice C is incorrect and may result from conceptual or calculation errors.

Question Difficulty: Easy

Question ID 9f0eba60

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: 9f0eba60

The function f is defined by $f(x) = 4x - 3$. What is the value of $f(10)$?

- A. -30
- B. 37
- C. 40
- D. 43

ID: 9f0eba60 Answer

Correct Answer: B

Rationale

Choice B is correct. It's given that the function f is defined by $f(x) = 4x - 3$. Substituting 10 for x in the given function yields $f(10) = 4(10) - 3$, which is equivalent to $f(10) = 40 - 3$, or $f(10) = 37$. Therefore, the value of $f(10)$ is 37 .

Choice A is incorrect and may result from conceptual or calculation errors.

Choice C is incorrect. This is the value of $f(10)$ for the function $f(x) = 4x$, not $f(x) = 4x - 3$.

Choice D is incorrect. This is the value of $f(10)$ for the function $f(x) = 4x + 3$, not $f(x) = 4x - 3$.

Question Difficulty: Easy

Question ID b3e6012b

Assessment	Test	Domain	Skill	Difficulty
PSAT/NMSQT & PSAT 10	Math	Algebra	Linear functions	<div><div></div><div></div><div></div></div>

ID: b3e6012b

To repair a refrigerator, a technician charges **\$60** per hour for labor plus **\$120** for parts. Which function f represents the total amount, in dollars, the technician will charge for this job if it takes x hours?

- A. $f(x) = x + 120$
- B. $f(x) = 60x$
- C. $f(x) = 60x + 120$
- D. $f(x) = 60x - 120$

ID: b3e6012b Answer

Correct Answer: C

Rationale

Choice C is correct. It’s given that the technician charges **\$60** per hour for labor. Therefore, if the job takes x hours, the technician will charge $\left(\frac{\$60}{1 \text{ hour}}\right)(x \text{ hours})$, or $\$60x$, for labor. It’s also given that the technician charges **\$120** for parts. Therefore, $f(x) = 60x + 120$ represents the total amount, in dollars, the technician will charge for this job if it takes x hours.

Choice A is incorrect and may result from conceptual or calculation errors.

Choice B is incorrect. This function represents the total amount, in dollars, the technician charges for labor only, not the total amount charged for labor and parts.

Choice D is incorrect. This function represents the total amount, in dollars, the technician would charge if the charge for parts were subtracted from, rather than added to, the charge for labor.

Question Difficulty: Easy